

## 2016 GPRO Quality Measures and Coding Checklist

Patient Name \_\_\_\_\_ DOB \_\_\_\_\_

Version 1.0 - 02052016

Measure and Description	Exclusion or Exceptions	ICD-10	CPT	CPT Description	Date Completed	Check if Billed
<p><b>Fall Screening</b> Patient considered at risk for future falls if they had 2 or more falls in the past years, or any fall with injury in the past year.</p> <p>*Screen patient 65 years or older for future fall risk at least once in measurement period</p>	Documentation of medical reason for not screening for fall risk (e.g., If patient is not ambulatory)	<p>Z91.81 At risk for falls, w/o falls</p> <p>Z78.9 At low risk for falls</p>	<p>1100F, 3288F</p> <p>1101F</p>	<p>Screened for future fall risk; documentation of 2 or more falls in past year or any fall w/ injury in past year; plan documented</p> <p>Screened for future fall risk; documentation of no falls in past year or only 1 fall w/o injury in past year; plan documented</p>		
<p><b>Clinical Depression</b> Screen for clinical depression and document follow-up if positive for patients 12 years or older.</p> <p>Administer PHQ-9 if answers on patient checklist are positive.</p> <p>Document plan of care on date of positive outcome</p> <p>*Screen at least once during measurement period</p>	<p>1) Patient already has active dx of depression or Bipolar disorder</p> <p>2) Patient refuses</p> <p>3) Situation is considered urgent or emergent</p> <p>4) If patients functional capacity or motivation to improve may impact the accuracy of results</p>	Z13.89 Encounter for screening for other disorder	G0444  G8431  G8510	<p>Annual depression screening performed (15 min) *bill with other code below at AWW</p> <p>Screen performed, clinically significant depressive symptoms, w/follow up plan</p> <p>Screen performed, negative for depression, no follow up required</p>		
<p><b>Depression Remission</b> If patients 18 years and older with major depression or dysthymia and had an initial PHQ-9 score &gt;9, they must demonstrate remission at 12 months (+/- 30 days) defined as PHQ-9 score &lt;5.</p> <p>This measure applies to both patients with newly diagnosed and existing depression whose current PHQ-9 score indicates a need for treatment</p>	<p>1) Patients who died</p> <p>2) Patients who received hospice or palliative care services</p> <p>3) Patients who were permanent nursing homes residents</p> <p>4) Patients who are diagnosed Bipolar Disorder</p> <p>5) Patients who are diagnosed with a personality disorder</p>	Z13.89 Encounter for screening for other disorder	3092F	Screening for clinical depression documented as positive, a follow-up plan not documented, documentation stating the patient is not eligible		
<p><b>Medication Documentation</b> Document a list of current meds on the date of encounter for patients 18 years or older *during all office visits;</p> <p>Must include all known Rx meds, OTC, herbals, and vitamins/minerals/dietary supplements and must contain name, dosage, frequency and route of administration;</p> <p>Must document if not currently taking any medications</p>	If situation is considered urgent or emergent		G8427	Medications reconciled in the medical record. All office visits during 12 month reporting period.		
<p><b>Body Mass Index Screening</b> For patients 18 years or older, calculate and documented BMI during current encounter or during the previous 6 months AND with a BMI out of normal parameters, a follow-up plan needs to be documented during the encounter or during the previous 6 months of the current encounter.</p> <p>**Age 65 + years -- BMI &gt; or = 23 and &lt;30 kg/m2 **Age 18 - 64 years -- BMI ≥18.5 and &lt;25 kg/m2</p> <p>*Screen at least once during measurement period</p>	<p>1) Patient is pregnant</p> <p>2) Patient refuses measurements</p> <p>3) Situation is urgent or emergent</p> <p>4) Provider documents why measurement was not appropriate</p> <p>5) Receiving Palliative care</p>		G8420  G8417  G8418	<p>BMI documented within normal limits, no follow-up required (23-29)</p> <p>BMI documented above normal, with follow-up documented (&gt; or = 30)</p> <p>BMI documented below normal, with follow-up documented (&lt;23)</p>		
<p><b>Screening for High Blood Pressure and Follow-up Plan</b> Screen for high blood pressure and document follow-up plan for patient 18 years or older if pre-hypertensive SBP 120-139 or DBP 80-89 or hypertensive BP &gt;140 or &gt;90.</p> <p>*Screen at least once during measurement period</p>	<p>1) Patient already has active dx of HTN</p> <p>2) Patient refuses either BP or follow up plan</p> <p>3) Situation is considered urgent or emergent</p>	<p>Z01.30 Exam of BP w/o abnormal findings</p> <p>Z01.31 Exam of BP w abnormal findings</p>	G8784  G8783  G8950	<p>Patient not eligible (e.g. due to active dx of HTN, patient refuses or urgent or emergent situation)</p> <p>Normal BP reading, no follow-up required</p> <p>Pre-hypertensive or hypertensive w/follow-up</p>		

Measure and Description	Exclusion or Exceptions	ICD-10	CPT	CPT Description	Date Completed	Check If Billed
<p><b>Controlling High Blood Pressure</b> BP is considered controlled in patients who have a diagnosis of HTN and whose BP is adequately controlled (&lt;140/90 mmHg) at the most recent visit during the measurement period for patients 18 to 85 years.</p> <p>BP readings performed by clinicians in the provider office are acceptable. However, BP reading by patient at their home are not acceptable.</p>	Pregnancy, ESRD, dialysis, or renal transplant before or during the measurement period	Z01.30 Exam of BP w/o abnormal findings  Z01.31 Exam of BP w abnormal findings	G8752  G8753  G8754  G8755	Most recent systolic BP<140mm/Hg  Most recent systolic BP≥140mm/Hg  Most recent diastolic BP<90mm/Hg  Most recent diastolic BP≥90mm		
<p><b>Tobacco Use Screening and Cessation</b> Screen for tobacco use in patients 18 years or older at least one or more times within 24 months AND give cessation counseling intervention and/or pharmacotherapy if identified as a tobacco user</p>	Documented reason for no screening, such as limited life expectancy or other medical reason	Z72.0 Tobacco user  Z78.9 Non tobacco user	G0436  G0437  G9458  G9459	Smoking cessation counseling greater than 3 minutes  Smoking cessation counseling greater than 10 minutes  Patient documented as a tobacco user and received cessation intervention  Currently a tobacco non-user		
<p><b>Influenza immunization</b> Patients 6 months or older seen for a visit between Oct 1, 2015 and March 31, 2016, should have influenza immunization OR who reported previous influenza immunization. Receipt of immunization from another provider or from same provider is required.</p> <p>*Vaccination once per year</p>	1) Medical reason for not receiving influenza immunization (e.g., allergy, intolerance) 2) Patient reason for not receiving influenza immunization (e.g., patient declined, other patient reasons) 3) System reason for not receiving influenza immunization (e.g., lack of drug availability, other reasons)	Z23 Encounter for immunization	G0008  G8482  G8483	Administration of influenza vaccine  Annual influenza vaccination administered or previously rec'd  Influenza vaccine not given (Declined or contraindicated for medical reasons)		
<p><b>Pneumonia Vaccination</b> Patients 65 years of age and older should receive one pneumococcal vaccine</p> <p>*Vaccination once in their lifetime</p>	None	Z23 Encounter for immunization	G0009  G8864  G8865  G8866	Administration of pneumococcal  Pneumonia vaccine administered or previously received at least 1 time after age 65  Pneumonia vaccine not administered for medical reasons and document  Pneumonia vaccination refusal		
<p><b>Diabetes - HbA1c control</b> Report most recent HbA1c level (performed during the measurement period). This measure is looking at the percentage of patients who have HbA1c &gt;9% in patients aged 18 to 75 years for patients with diabetes type 1 or type 2.</p> <p>*Report most recent HbA1c at visit during measurement</p>	None	Z13.1 Encounter for screening for diabetes	3044F  3045F  3046F	A1c<7%  A1c 7-9%  A1c>9%		
<p><b>Diabetes - Eye Exam</b> Patient must have retinal or dilated eye exam by eye care professional *1 time during measurement period or a negative retinal exam (no evidence of retinopathy) in the 12 months prior to measurement period in patients age 18 to 75 years for patients with diabetes type 1 or type 2.</p>	None	Z13.5 Screen for Eye disorder	2022F  3072F	Dilated eye exam w/ interpretation by oph or optom. Documented and reviewed  Low risk for retinopathy		
<p><b>Breast Cancer Screening</b> Women 50 - 74 years of age with mammogram to screen for breast cancer within 27 months; document results obtained and discussed with patient</p>	Women who have had a bilateral mastectomy or for whom there is evidence of two unilateral mastectomies	Z12.31 Encounter for mammogram screening	G0202	Screening mammogram		

Measure and Description	Exclusion or Exceptions	ICD-10	CPT	CPT Description	Date Completed	Check If Billed
<b>Colorectal Cancer Screening</b> Adults 50 - 75 years of age with appropriate screening for colorectal screening as defined by - Fecal occult blood test (FOBT) 1 time during measurement period  - Flex sigmoidoscopy during measurement period or the 4 years prior to measurement period  - Colonoscopy during measurement period or the 9 years prior to the measurement period	Patients with diagnosis of past history of total colectomy or colorectal cancer	Z12.11 Encounter for screening for cancer of colon	G0328 G0104 G0105 G0464	Fecal Occult blood test, immunoassay  Colorectal Cancer screening; flex sig  Colorectal Cancer screening; colonoscopy  Colorectal cancer screening;stool-based DNA and fecal occult hemoglobin (e.g., KRAS, NDRG4 and BMP3)		
<b>Coronary Artery Disease</b> Patients 18 years or older, with a diagnosis of:  - CAD with Diabetes OR - CAD and a current or prior LVEF <40%  then ACE or ARB must be prescribed	1) Medical reason for not prescribing (e.g., allergy, intolerance) 2) Patient reason for not prescribing (e.g., patient declined, other patient reasons) 3) System reason for not prescribing (e.g., lack of drug availability, other reasons)	I25.1 Atherosclerotic cardiovascular disease  E10 Type 1 DM  E11 Type 2 DM  I50.9 Heart Failure, unspecified	4010F	ACE or ARB therapy prescribed or taken		
<b>Heart Failure</b> Patients 18 years or older with: - current HF diagnosis and with a current or prior LVEF <40%  then BB therapy must be prescribed either within a 12 month period when seen in outpatient setting OR at each hospital discharge.	1) Medical reason for not prescribing (e.g., allergy, intolerance) 2) Patient reason for not prescribing (e.g., patient declined, other patient reasons) 3) System reason for not prescribing (e.g., lack of drug availability, other reasons)	I50.9 Heart Failure, unspecified	4008F  G8395	Beta blocker therapy prescribed or currently taking  LVEF >= 40% or documentation as normal or mildly depressed LVEF		
<b>Ischemic Vascular Disease</b> Patients 18 years or older: - who have been discharged for AMI, CABG, or PCI in the 12 months prior to the measurement period, OR - who have an active diagnosis or IVD during the measurement period,  should have documentation of use of ASA or another antithrombotic	None	Z79.82 Long term ASA use	G8598	ASA or another antithrombotic therapy prescribed/used		
<b>Statin Therapy for CV Disease</b>  - Adults aged 21 years or older who were previously diagnosed with/or currently have an active diagnosis of clinical atherosclerotic cardiovascular disease (ASCVD); OR - Adults aged 21 years or older with a fasting or direct low-density lipoprotein cholesterol (LDL-C) level >190mg/dL; OR - Adults aged 40-75 with a diagnosis of diabetes with a fasting or direct LDL-C level of 70-189mg/dL  should be prescribed statin therapy.	1) Patient with adverse effect, allergy, intolerance to statin 2) Pregnant or breastfeeding 3) Receiving Palliative care 4) Active liver disease or hepatic disease or insufficiency 5) ESRD 6) Patient with diabetes, fasting or direct LDL-C <70 mg/dL and on statin	I25.1 Atherosclerotic cardiovascular disease	G9662  G9667  G9664  3048F  3049F  3050F	Previously dx or have active diagnosis of clinical ASCVD  Documentation of medical reason for not being on statin therapy  Currently statin therapy or received Rx for statin  Most recent LDL-C less than 100mg/dL  Most recent LDL-C 100-129 mg/dL  Most recent LDL-C greater than or equal to 130 mg/dL		